


Section 9.1. The Standard Atmosphere



Atmospheric Model Background

- Earth's atmosphere is an extremely thin sheet of air extending from surface of Earth to edge of space.
- If Earth were size of a basketball, a tightly held pillowcase would represent thickness of atmosphere.
- Gravity holds atmosphere to Earth's surface. Within atmosphere, very complex chemical, thermodynamic, and fluid dynamics effects occur.
- Atmosphere is not uniform; fluid properties are constantly changing with time and place. We call this change “weather”.
- Variations in air properties extend upward from surface of Earth. Sun heats surface of Earth, and some of this heat goes into warming air near surface, heated air is also diffused or convected up through atmosphere.

Atmospheric Model Background (2)

- Thus the air temperature (and the local speed of sound) is highest near the surface and decreases as altitude increases.
- Air pressure is proportional to the weight of the air over a given location. Thus air pressure decreases as we increase altitude.
- Air density depends on both the temperature and the pressure through the equation of state and also decreases with increasing altitude.
- Due to localized sun angles, terrain, gravitational changes with latitude, and coriolis forces due to earth's rotation, properties within the atmosphere change significantly with time of year, and global latitude/longitude.
- To help with establishing “mean designs” for air-vehicles, it is extremely useful to define “standard” conditions, and develop models of how the atmospheric properties vary as a function of altitude .. Thus the so-called Standard Day, and Standard Atmosphere.”

Hydrostatic Model of the Atmosphere

- For a fluid in rest without shearing stresses, any elementary fluid element will be subjected to two types of forces, namely, surface forces due to the pressure and a body force equal to the weight of the element.

Hydrostatic Equation


$$\frac{\partial p}{\partial z} = -\rho \cdot g = -\frac{p}{R_g \cdot T} \cdot g$$

Integrating over a depth of atmosphere

Assuming constant gravity

$$\int_{p_1}^{p_2} \frac{dp}{p} = \ln\left(\frac{p_2}{p_1}\right) = -\frac{g}{R_g} \int_{z_1}^{z_2} \frac{d\xi}{T(\xi)}$$

$$T(z) = T_1 - \alpha_L \cdot z \rightarrow \alpha_L \equiv \text{Lapse Rate}$$



Need temperature profile
To complete integral

Temperature "Lapse Rate" Model

Hydrostatic Model of the Atmosphere (2)

$$\ln\left(\frac{p_2}{p_1}\right) = -\frac{g}{R_g} \int_{z_1}^{z_2} \frac{d\xi}{T(\xi)} = -\frac{g}{R_g} \int_{z_1}^{z_2} \frac{d\xi}{T_1 - \alpha_L \cdot \xi}$$

$$T(z) = T_1 - \alpha_L \cdot z \rightarrow \alpha$$

$$\ln\left(\frac{p_2}{p_1}\right) = -\frac{g}{R_g} \int_{z_1}^{z_2} \frac{d\xi}{T_1 - \alpha_L \cdot \xi} =$$

$$\frac{g}{\alpha_L R_g} \left[\ln(T_1 - \alpha_L \cdot z_2) - \ln(T_1 - \alpha_L \cdot z_1) \right] =$$

$$\frac{g}{\alpha_L R_g} \ln\left(\frac{T_1 - \alpha_L \cdot z_2}{T_1 - \alpha_L \cdot z_1}\right)$$

Hydrostatic Model of the Atmosphere (3)

Let $\rightarrow z_1 = 0, T = T_{sl}$

$$\rightarrow \ln\left(\frac{p_2}{p_{SL}}\right) = \frac{g}{\alpha_L R_g} \ln\left(\frac{T_1 - \alpha_L \cdot z_2}{T_1}\right) = \frac{g \cdot T_{sl}}{\alpha_L R_g} \ln\left(1 - \frac{\alpha_L}{T_{sl}} \cdot z_2\right)$$


$$p(z) = p_{sl} \cdot \left(1 - \frac{\alpha_L}{T_{sl}} \cdot z\right)^{\frac{g}{\alpha_L R_g}}$$

Decay of Pressure with Altitude in Troposphere

Hydrostatic Model of the Atmosphere (4)


- For the stratosphere atmospheric layer (between 11.0 and 20.1 km), the temperature has a constant value (isothermal conditions) which is 56.5 C (or 69.7 F, 389.97 R, 216.65 K).

$$p(z) = p_s \cdot e^{-\frac{g}{R_g \cdot T_s} (z - z_s)}$$



Decay of Pressure with Altitude in Stratosphere

Extension to Mesosphere



International Standard Atmospheric Model


- There are a myriad of atmospheres available
 - .. Each serving a certain technical purpose ...

see link on section 4 wep page ..

http://www.spacewx.com/Docs/AIAA_G-003B-2004.pdf

- Basically, working models break into distinct altitude layers

- 1) Troposphere
 - $\rightarrow h < 11 \text{ km (36,000 ft)}$
- 2) Lower Stratosphere
 - $\rightarrow 11 \text{ km} \leq h < 25 \text{ km (82,000 ft)}$
- 3) Upper Stratosphere
 - $\rightarrow 25 \text{ km} \leq h < 50 \text{ km}$
- 4) Mesosphere
 - $\rightarrow h > 50 \text{ km}$




International Standard Atmospheric Model (2)

| Layer | Level Name | Base Geopotential Altitude above MSL ^[5] <i>h</i> (m) | Base Geometric Altitude above MSL ^[5] <i>z</i> (m) | Lapse Rate (°C/km) ^[a] | Base Temperature <i>T</i> (°C) | Base Atmospheric Pressure <i>p</i> (Pa) | Base Atmospheric Density <i>ρ</i> (kg/m ³) |
|-------|--------------|---|--|--------------------------------------|-----------------------------------|--|---|
| 0 | Troposphere | -610 | -611 | -6.5 | +19.0 | 108,900 (1.075 bar) | 1.2985 |
| 1 | Tropopause | 11,000 | 11,019 | +0.0 | -56.5 | 22,632 | 0.3639 |
| 2 | Stratosphere | 20,000 | 20,063 | +1.0 | -56.5 | 5474.9 | 0.0880 |
| 3 | Stratosphere | 32,000 | 32,162 | +2.8 | -44.5 | 868.02 | 0.0132 |
| 4 | Stratopause | 47,000 | 47,350 | +0.0 | -2.5 | 110.91 | 0.0020 |
| 5 | Mesosphere | 51,000 | 51,413 | -2.8 | -2.5 | 66.939 | |
| 6 | Mesosphere | 71,000 | 71,802 | -2.0 | -58.5 | 3.9564 | |
| 7 | Mesopause | 84,852 | 86,000 | - | -86.28 | 0.3734 | |

^a lapse rate given per kilometer of *geopotential altitude*

In the above table, *geopotential altitude* is calculated from a mathematical model that adjusts the altitude to include the variation of gravity with height, while *geometric altitude* is the standard direct vertical distance above mean sea level.^[2] Note that the Lapse Rates cited in the table are given as °C per kilometer of geopotential altitude, not geometric altitude.

1976 US Standard Atmospheric Model



1976 US Standard Atmospheric Model (2)

$$\frac{z}{H_s} \rightarrow \begin{aligned} z &= \text{geometric altitude} \\ H_s &= \text{"scale height"} \end{aligned}$$

1976 US Standard Atmosphere

$$H_s = \frac{c_0^2}{\gamma \cdot g_0} = \frac{R_g \cdot T_0}{g_0}$$

$$T_{SL} = 518.67^{\circ}R (59^{\circ}F)$$

$$T_{SL} = 288.15^{\circ}K (15^{\circ}C)$$

$$P_{SL} = 2116.22 \text{ pounds/ft}^2$$

$$P_{SL} = 1.013250 \times 10^5 \text{ newtons/m}^2$$

$$\rho_{SL} = 0.0023769 \text{ slug/ft}^3$$

$$\rho_{SL} = 1.2250 \text{ kilograms/m}^3$$

$$g = 32.1741 \text{ ft/sec}^2$$

$$g = 9.80665 \text{ m/sec}^2$$

$$a_0 = 1116.45 \text{ ft/sec}$$

$$a_0 = 340.294 \text{ m/sec}$$

$$H = 27,672 \text{ ft}$$

$$H = 8434.5 \text{ m/sec}$$

$$R_{air} = 1710.2 \text{ ft}^2 / (\text{sec}^2 - {}^{\circ}R)$$

$$R_{air} = 287.06 \text{ m}^2 / (\text{sec}^2 - {}^{\circ}K)$$

1976 US Standard Atmospheric Model (3)

**Earth Atmosphere Model**
*Metric Units*Glenn
Research
CenterFor $h > 25000$ (Upper Stratosphere)

$$T = -131.21 + .00299 h$$

$$p = 2.488 * \left[\frac{T + 273.1}{216.6} \right]^{-11.388}$$

For $11000 < h < 25000$ (Lower Stratosphere)

$$T = -56.46$$

$$p = 22.65 * e^{(1.73 - .000157 h)}$$

For $h < 11000$ (Troposphere)

$$T = 15.04 - .00649 h$$

$$p = 101.29 * \left[\frac{T + 273.1}{288.08} \right]^{5.256}$$


1976 US
Standard
AtmosphereUp Through
Stratosphere ρ = density (kg/cu m) p = pressure (K-Pa)

$$\rho = p / (.2869 * (T + 273.1))$$


 T = temperature ($^{\circ}$ C) h = altitude (m)

1976 US Standard Atmospheric Model(4)

Temperature / altitude graph



Pressure / altitude graph



1976 US Standard Atmosphere up Through Mesosphere

1976 US Standard Atmospheric Model(5)

| Altitude (m) | Temperature (K) | Pressure ratio | Density (kg/m ³) |
|---------------|-----------------|--------------------------|---------------------------------|
| 0 (sea level) | 288.150 | 1.0000 | 1.2250 |
| 1000 | 281.651 | 8.87×10^{-1} | 1.11117 |
| 3000 | 268.650 | 6.6919×10^{-1} | 0.90912 |
| 5000 | 255.65 | 5.3313×10^{-1} | 0.76312 |
| 10,000 | 223.252 | 2.6151×10^{-1} | 4.1351×10^{-1} |
| 25,000 | 221.552 | 2.5158×10^{-2} | 4.0084×10^{-2} |
| 50,000 | 270.650 | 7.8735×10^{-4} | 1.0269×10^{-3} |
| 75,000 | 206.650 | 2.0408×10^{-5} | 3.4861×10^{-5} |
| 100,000 | 195.08 | 3.1593×10^{-7} | 5.604×10^{-7} |
| 130,000 | 469.27 | 1.2341×10^{-8} | 8.152×10^{-9} |
| 160,000 | 696.29 | 2.9997×10^{-9} | 1.233×10^{-9} |
| 200,000 | 845.56 | 8.3628×10^{-10} | 2.541×10^{-10} |
| 300,000 | 976.01 | 8.6557×10^{-11} | 1.916×10^{-11} |
| 400,000 | 995.83 | 1.4328×10^{-11} | 2.803×10^{-12} |
| 600,000 | 999.85 | 8.1056×10^{-13} | 2.137×10^{-13} |
| 1,000,000 | 1000.00 | 7.4155×10^{-14} | 3.561×10^{-15} |

1976 US Standard Atmosphere Source Code

1976 US Standard Atmosphere up Through Mesosphere

Earth Global Reference Atmospheric Model (Earth-GRAM 2010)

GRAM ..

- Global model, using empirical database, giving density, temperature, pressure, winds, and selected atmospheric constituent concentrations, from the surface of the Earth to orbital altitudes, as a function of geographic position and time of year.
- Local perturbations about mean conditions are also included. Model is also suitable for use as a subroutine in a trajectory code or orbit propagator program or other programs used for simulations of in-flight or on-orbit atmospheric variability in density, temperature, or winds.

<https://see.msfc.nasa.gov/model-gram>

Questions??

